EVOLUTION

INTERNATIONAL JOURNAL OF ORGANIC EVOLUTION

Volume II, 1948

Published Quarterly

At Prince and Lemon Streets, Lancaster, Pa.

By the

Society for the Study of Evolution

TABLE OF CONTENTS

Number 1, March

Anderson, Edgar. Hybridization of the habitat 1
Shull, A. Franklin. Natural hybrids of subspecies of *Hippodamia quinquesignata* (Coccinellidae) 10
Lerner, I. Michael, and Everett R. Dempster. Some aspects of evolutionary theory in the light of recent work on animal breeding 19
Lutz, Bertha. Ontogenetic evolution in frogs 29
Reed, S. C., and E. W. Reed. Morphological differences and problems of speciation in *Drosophila* 40
Hsu, T. C., and T. T. Liu. Microgeographic analysis of chromosomal variation in a Chinese species of *Chironomus* (Diptera) 49
Pittendrigh, Colin S. The Bromeliad-Anopheles-Malaria complex in Trinidad 58
White, M. J. D. The chromosomes of the parthenogenetic mantid *Brunneria borealis* 90

Number 2, June

Lack, David. Natural selection and family size in the starling 95
Chapin, James P. Variation and hybridization among the Paradise Flycatchers of Africa 111
Axelrod, Daniel I. Climate and evolution in western North America during Middle Pliocene time 127
Colbert, Edwin H. Evolution of the Horned Dinosaurs 145
Schaeffer, Bobb. The origin of a mammalian ordinal character 164
Reed, S. C., and E. W. Reed. Natural selection in laboratory populations of *Drosophila* 176
Streisinger, George. Experiments on sexual isolation in *Drosophila*. IX. Behavior of males with etherized females 187

Number 3, September

Wallace, Bruce. Studies on "sex-ratio" in *Drosophila pseudoobscura*. I. Selection and "sex-ratio" 189
Rensch, Bernhard. Histological changes correlated with evolutionary changes of body size 218
Lundman, Bertil. Geography of human blood groups (A, B, O system) .. 231
Mayr, Ernst, and Charles Vaurie. Evolution in the family Dicruridae .. 238
Hairston, Nelson G., and Clifford H. Pope. Geographic variation and speciation in Appalachian Salamanders (*Plethodon jordani* group) 266

Number 4, December

Wright, Sewall. On the roles of directed and random changes in gene frequency in the genetics of populations 279
Stalker, Harrison D., and Hampton L. Carson. An altitudinal transect of *Drosophila robusta* Sturtevant 295

GOSLINE, WILLIAM A. Speciation in the fishes of the genus *Menidia* 306
RAND, A. L. Glaciation, an isolating factor in speciation 314
CLOUD, PRESTON E. JR. Some problems and patterns of evolution exemplified by fossil invertebrates 322
ZANGERL, RAINER. The methods of comparative anatomy and its contribution to the study of evolution 351
MAYR, E. Climatic races in plants and animals 375

INDEX TO VOLUME II, 1948

acceleration, 340
Achillea, 375–376
adaptive, convergence, 327; radiation, 146; values, 211
Africa, West, 118
alleles, blood, 231
Allen's rule, 255
allometry, in embryology, 218–230; in evolution, 218–230
altitudinal transect, 295–305
Ammonoid cephalopods, 329
Anchiceratops, 145
Anderson, Edgar, 1–9
animals, pre-Cambrian, 347
Anopheles, 58; *A. bellator*, 58; *A. homunculus*, 59
antelopes, 162
Arrhinoceratops, 145
Artiodactyla, 164
astragalus, 167
Axelrod, Daniel I., 127–144
back-crosses, 1–9
banding, in *Hippodamia*, 16
behavior, *Drosophila* males with etherized females, 187–188
Bergmann's rule, 253
blood allele, in Europe (map of), 234; *p* (map of), 232; *q* (map of), 233; *r* (map of), 235
blood groups, human, 231–237
body size, evolutionary changes of, 218–230; lower limit, of, 227; increase of, 229
borderline, between species and subspecies, 261
Brachiopods, 323, 330, 332; terebratuloid, 343
brain, 228; modification of structure, 230
Branta canadensis, 315
breeding, animal, 19–28
Bromeliad-Anopheles-Malaria complex in Trinidad, 58–89
Bromeliads, 32, 58–89
Bromelioideae, 64, 85
Brunneria borealis, 90–93
Bufo paracnemis, 35
Cambrian fauna, origin of, 346
Canachites canadensis, 317; *C. franklini*, 317
carnivores, 172
Carson, Hampton L., 295–305
categories, origin of higher, 241
Centrolenella eurygnatha, 31, 33
Ceratopsia, 145–163, 152; evolution of, 160; frill, 150; distribution of (table), 160; nasal horn, 150
Chaetorhynchus, 241
Chapin, James, 111–126
characters, primitive, 275; taxonomic, 241
Chasmosaurus, 145
Cheirothyris, 323
Chironomus, 49–57
chromosomes, 90–93, 190; salivary gland, 56; X, 189
Clausen, J., 375
climate, 62; and evolution, 127–144
climatic races, 375–376
clines, 252, 254, 257, 308, 375; altitudinal, 295; progressive, 257
clisere, 131
Cloud, Preston, E. Jr., 322–350
clutch-size, 96
Coccinellidae, 10–18
coefficients, path, 19
Coelonotus fissilis, 33
Colaptes auratus, 111; *C. cafer*, 111
Colbert, Edwin H., 145–163
comparative morphology, methods of, 354
competition, 345; interspecific, 31, 33; larval, 198
convergence, 324; adaptive, 327
Cope's rule, 218, 228
Corvus corone, 111; *C. cornix*, 111
creodonts, 172
Cretaceous, 153
curves, survivorship, 202
Cyclorhamphus fuliginosus, 32; *C. pinderi*, 32
Dempster, Everett R., 19–28
Dendroica auduboni, 318; *D. coronata*, 318
Dermochelys, 359
development, 30; acceleration, 35
Dicruridae, 238–265; genera of, 241; *D. hottentottus*, 250; *D. paradiseus* (map of), 251
differences, morphological, 40–48; species, 42
dimorphism, sexual, 271
dinosaurs, horned, 145–163; evolution of, 145–163
discriminant function, 301
dispersal, 263
Dobzhansky, Th., 43
drift, genetic, 183
drongos, 238–265; parallelism in, 242; size variation in, 242
Drosophila, 40–48, 189–217; natural selection in laboratory populations, 176–186; sex-ratio and selection and sex-ratio in, 189–217; *D. melanogaster*, 176, 187; *D. miranda*, 41; *D. persimilis*, 41, 187, 194, 218; *D. pseudoobscura*, 41, 187, 298; *D. subobscura*, 41; *D. robusta*, 295–305
ecotype, 375
Edwardsia, 337
eggs, hatchability, 210; limitation of number, 227
Eleutherodactylus, 37; *E. nasutus*, 33; *E. guentheri*, 33
embryology, allometry in, 218–230
environment, 298
equilibrium, 56, 177, 292, 375

eruptive evolution, 342
Erythrina micropteryx, 62
evolution, allometry in, 218–230; and climate, 127–144; and morphology, 371; ceratopsian, 160; during Middle Pliocene, 133; eruptive, 342; in family Dicruridae, 238–265, mega-, 171; of Horned Dinosaurs, 145–163; ontogenetic, 29–39; parallelism in, 246; quantum, 171; tachytelic, 171
evolutionary change, rate of, 171
evolutionary rates, 152; theory, 19–28
fecundity, 204
fitness, 21
flora, Bromeliad, 58–89; middle Pliocene, 133
fluctuations, in gene frequency, 282; in selective value, 285
flycatchers, Paradise, 111–126
frequencies, gene, 281
Gastropods, 329
Gavia adamsii, 317; *G. immer*, 317
gene, additive action, 25; arrangement, 54, 176, 297; frequencies, 281; recessive lethal, 190
genetic drift, 183
genetics, population, 279
genotype, aggregate, 21
geographical isolation, 313
geographical variation, 259–306; in Appalachian Salamanders, 266–278; of habitat requirements, 258
glaciation, 314–321
Gosline, William A., 306–313
growth, allometric, 218
habitat, 320; disturbance and hybridization, 111–126; hybridization of, 1–9
Hairston, Nelson G., 266–278
Haldane, J. B. S., 21
hatchability, egg, 210
heritability, 20
heterozygosis, 14
heterozygote, superiority of, 181, 214
Hexacorallia, 334
Hiesey, W. M., 375
Hippodamia, banding in, 16; spotting in, 15; *H. sinuata spuria*, 10–18; *H. quinquesignata*, 10–18
homology, 351, 362; definition of, 366
homeomorphy, 323
Hsu, T. C., 49–57
Huxley, J., 38
hybrid swarms, 111
hybrids, 10–18, 311
hybridism, interspecific, 117
hybridization, 111–126; introgressive, 3, 111; of the habitat, 1–9
Hyla goeldii, 33; *H. mesophaea*, 33; *H. ohausi*, 32; *H. polytaenia*, 33
Hyopsodontidae, 166
immigration, 293
index, selection, 22; tail, 247; wing, 42
insects, nervous system, 223
introgressive hybridization, 3, 111
invasions, double, 262
inversions, heterozygote, 176; tandem, 55; terminal, 190
invertebrates, fossil, 322–350
Iris fulva, 6; *I. giganticaerulea*, 6
isoalleles, 289
isolation, geographic, 276, 314; sexual in *Drosophila*, 187–188
Junco hyemalis, 318; *J. oreganus*, 318
Keck, D. D., 375
Kerteszia, 58–89
laboratory populations of *Drosophila*, 176–186
Lack, D., 95–110
Larus argentatus, 315; *L. leucopterus*, 315
Leptoceratops, 145
Leptodactylus mystaceus, 32; *L. nanus*, 33
Lerner, I. M., 19–28
light, effect of, 70
Liu, T. T., 49–57
longevity of *Drosophila* adults, 201
Lundman, Bertil, 231–237
Lutz, Bertha, 29–39
mammals, orders of, 164
mantid, 90–93
Mather, K., 43
Mayr, E., 238–265, 375–376
Menidia, 306–313; *M. audens*, 306; *M. beryllina*, 306; *M. extensa*, 306; *M. menidia*, 306–313
methods of comparative anatomy, 351–374
microgeographic analysis, 49–57
mites, 191
modifiers, 289
Monoclonius, 145
morphological concepts, 351; differences, 40–48; series, 369
morphology, comparative, 351; and evolution, 371
morphotype, 355
mortality, selective, 198
mutations, pressure, 281; randomness of, 246
Narcissus, 2
natural hybrids of subspecies of *Hippodamia quinquesignata* (*Coccinellidae*), 10–18
natural selection, 24, 215, 218; and family size in starling, 95–110; in laboratory populations of *Drosophila*, 176–186
neoteny, 331
nervous system, in insects, 223
northernness, 303
Omphalophyllia, 340
ontogenetic evolution, 29–39
origin of a mammalian ordinal character, 164–175
Ornithischia, 145
Paedomorphosis, 331
paleobotany, 127–144
Paleocene, 171
Panaxia dominula, 281
parallelism, 304, 324; in drongos, 242
parthenogenesis, 90
Parus cinctus, 317; *P. hudsonicus*, 317
Pelecypods, 331

Pentaceratops, 145
Phyla, 322
Phyllomedusa guttata, 31, 33
phylogenetic trees, 239
phylogeny, 342
Physalaemus fuscomaculatus, 33
Pitcairnioideae, 63, 85
Pittendrigh, C. S., 58–89
Plethodon jordani, 266–278
Plethodontidae, 266
Pliocene, 127–144; floras, 133
polymorphism, 113; balanced, 176
Pope, C. H., 266–278
populations, 306, 375; bottle, 177; cages, 176, 191; experimental, 198; genetics, 279; hybrid, 262; laboratory, 176–186; on small islands, 255; peripheral, 260; size, 283, 284; small isolated, 171; smallness of, 279; structure, 280; turn-over, 106
potentialities, evolutionary, 246
preference, sexual, 209
problems and patterns of evolution exemplified by fossil invertebrates, 322–350
proportions, 244, 247
Protoceratops, 145
Pycnoscelus surinamensis, 91
race, climatic, 375–376; ecological, 375
radiation, adaptive, 146
Rand, A. L., 314–321
random fluctuations, 286
randomness, 246
rate, evolutionary, 152, 171
Reed, E. W., 40–48, 176–186
Reed, S. C., 40–48, 176–186
refugia, 314
relationships, intrageneric, 275
Rensch, B., 218–230
roles of directed and random changes in gene frequency in genetics of populations, 279–294
rules, ecological, 252
Saga pedo, 91
salamanders, Appalachian, 266–278; speciation in, 266–278
salivary gland chromosome, 56
sampling, accidents of, 281
Schaeffer, Bobb, 164–175
selection, and family size in starling, 95–110; and sex-ratio, 189–217; index, 22; natural, 24, 95–110; pressure, 21, 30, 172, 292, 345, 375; sequential, 25
selective, forces, 198; mortality, 198; value, 184
semi-species, 314
serological regions (map of), 236
sex-comb, 42
sex-ratio, 189–217; and selection, 189–217; geographical distribution, 191 (map of), 192; in *Drosophila pseudoobscura*, 189–217, 194
sexual, activity, 207; isolation in *Drosophila*, 187–188; preference, 209
Shull, A. F., 10–18
size, family, 95–110; clutch, 96; gradient, 271; variation on islands, 253
skull, 356; of turtles, 359
Sophophora, 189
southernness, 303
speciation, 258, 314; by geographic isolation, 276; in Appalachian salamanders, 266–279; in fishes of genus *Menidia*, 306–313
species, character, 247; differences, 42; primitive, 239; specialized, 239
Stalker, H. D., 295–305
starling, 95–110
sterility, 1
Streisinger, G., 187–188
Stropheodontidae, 325
Strophomenidae, 325
structural plan, 355
structure, histological, 220; modification of brain, 230
Sturnus vulgaris, 96
Styracosaurus, 145
subspeciation, rate of, 259
subspecies, number of, 258
superspecies, 259, 261
survival, 99
survivorship curves, 202
tarsal joint, 169
Terpsiphone, 111–126; *T. rufiventer*, 113 (map of), 115, 120; *T. rufiventer somereni*, 114 (map of), 115, 120; *T. rufocinerea*, 113 (map of), 116; *T. viridis*, 113 (map of), 116; *T. rufocinerea* x *viridis*, 118; *T. rufiventer* x *rufocinerea*, 117; *T. viridis* x *rufiventer*, 118, 119
Tetracorallia, 331, 336
Tetractinella, 323
Theophyllum splitgerberi, 60
Theobroma cacao, 62
Tillandsioideae, 64, 85
Torosaurus, 145
Tradescantia, 2; *T. canaliculata*, 5; *T. subaspera*, 5
transect, altitudinal, 295–305
trends, evolutionary, 241
Triceratops, 145
Trilobita, 334
Trinidad, 58–89
turn-over, population, 106
turtle skull, 359
variance, 21
variation, 111–126; chromosomal, 49–57; correlated, 246; geographic, 113, 247, 251, 259, 266–278; independent, 246
Vaurie, C., 238–265
vegetation, types, 62
vertebrates, 225; brain of, 225
Wallace, B., 189–217
White, M. J. D., 90–93
wing, 228; index, 42
Wright, S., 19, 279–294
X-chromosome, 189
Zangerl, R., 351–374
Zonotrichia leucophrys, 318